

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 37/18	A1	(11) International Publication Number: WO 99/40788 (43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/US99/02554 (22) International Filing Date: 5 February 1999 (05.02.99) (30) Priority Data: 60/075,122 13 February 1998 (13.02.98) US (71) Applicant (for all designated States except US): AMYLIN PHARMACEUTICALS, INC. [US/US]; 9373 Towne Centre Drive, San Diego, CA 92121 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): YOUNG, Andrew, A. [US/US]; 9514 Easter Way, San Diego, CA 92121 (US). VINE, Will [US/US]; 14537 Crestline Drive, Poway, CA 92064 (US). BEELEY, Nigel, R., A. [US/US]; 227 Loma Corta Drive, Solana Beach, CA 92075 (US). PRICKETT, Kathryn [US/US]; 7612 Trailbrush Terrace, San Diego, CA 92126 (US). (74) Agents: MUNSON, Peter, R. et al.; Lyon & Lyon LLP, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071-2066 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: INOTROPIC AND DIURETIC EFFECTS OF EXENDIN AND GLP-1 (57) Abstract <p>Methods for increasing urine flow are disclosed, comprising administration of an effective amount of GLP-1, an exendin, or an exendin or GLP-1 agonist. Methods for increasing urinary sodium excretion and decreasing urinary potassium concentration are also disclosed. The methods are useful for treating conditions or disorders associated with toxic hypervolemia, such as renal failure, congestive heart failure, nephrotic syndrome, cirrhosis, pulmonary edema, and hypertension. The present invention also relates to methods for inducing an inotropic response comprising administration of an effective amount of GLP-1, an exendin, or an exendin or GLP-1 agonist. These methods are useful for treating conditions or disorders that can be alleviated by an increase in cardiac contractility such as congestive heart failure. Pharmaceutical compositions for use in the methods of the invention are also disclosed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DESCRIPTION

INOTROPIC AND DIURETIC EFFECTS OF EXENDIN AND GLP-1

5 FIELD OF THE INVENTION

The present invention relates to methods for increasing urine flow comprising administration of an effective amount of glucagon-like peptide-1[7-36] amide (abbreviated "GLP-[7-36]NH₂" or simply "GLP-1"), an exendin, or an exendin or GLP-1 agonist.

10 Methods for increasing urinary sodium excretion and decreasing urinary potassium concentration are also disclosed. The methods are useful for treating conditions or disorders associated with toxic hypervolemia, such as renal failure, congestive heart failure, nephrotic syndrome, cirrhosis, pulmonary edema, and hypertension.

Pharmaceutical compositions for use in the methods of the invention are also disclosed.

The present invention also relates to methods for inducing an inotropic response
15 comprising administration of an effective amount of an exendin, GLP-1, or an exendin or GLP-1 agonist. These methods are useful for treating conditions or disorders that can be alleviated by an increase in cardiac contractility, such as congestive heart failure.

The following description summarizes information relevant to the present invention. It is not an admission that any of the information provided herein is prior art to
20 the presently claimed invention, nor that any of the publications specifically or implicitly referenced are prior art to that invention.

GLP-1

Glucagon-like peptide-1 [7-36] amide (also referred to as GLP-1[7-36]NH₂ or
25 GLP-1) is a product of the proglucagon gene. It is secreted into plasma mainly from the gut and produces a variety of biological effects related to pancreatic and gastrointestinal function. The parent peptide, proglucagon (PG), has numerous cleavage sites that produce other peptide products dependent on the tissue of origin including glucagon (PG[32-62]) and GLP-1[7-36]NH₂ (PG[72-107]) in the pancreas, and GLP-1[7-37] (PG[78-108]) and
30 GLP-1[7-36]NH₂ (PG [78-107]) in the L cells of the intestine where GLP-1[7-36]NH₂ (78-107 PG) is the major product.

GLP-1[7-36]NH₂, also known as proglucagon [78-107], or commonly, just GLP-1," as used herein, has an insulinotropic effect, stimulating insulin secretion from pancreatic β -cells; GLP-1 also inhibits glucagon secretion from pancreatic α -cells (Orskov, et al., Diabetes, 42:658-61, 1993; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996).

5 GLP-1 is reported to inhibit gastric emptying (Williams B, et al., J Clin Endocrinol Metab 81 (1): 327-32, 1996; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993), and gastric acid secretion. (Schjoldager BT, et al., Dig Dis Sci 34 (5): 703-8, 1989; O'Halloran DJ, et al., J Endocrinol 126 (1): 169-73, 1990; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993). A diuretic, antidysogenic effect of intracerebroventricular administration of GLP-

10 1 has been reported, however, this report claims that a peripheral, intraperitoneal injection of GLP-1 did not have this effect. (Tand-Christensen et al., Am. J. Physiol., 271:R848-56, 1996). GLP-1[7-37], which has an additional glycine residue at its carboxy terminus, also stimulates insulin secretion in humans (Orskov, et al., Diabetes, 42:658-61, 1993). A transmembrane G-protein adenylate-cyclase-coupled receptor believed to be responsible

15 for the insulinotropic effect of GLP-1 has been cloned from a β -cell line (Thorens, Proc. Natl. Acad. Sci., USA 89:8641-45, 1992).

Glucagon and glucagon-like peptides have been found to have different cardiovascular effects. Glucagon has been reported to have positive inotropic and chronotropic effects, produce a slight increase in arterial blood pressure in normal

20 individuals, and affect regional blood circulation. GLP-1 has been found to produce a moderate increase in both systolic and diastolic blood pressure, while GLP-2 has no effect on those parameters. GLP-1, administered through the jugular vein, has been reported to induce an increase in systolic and diastolic blood pressure and heart rate. (Reviewed in Barragán, J.M., et al., Regul. Peptides, 67:63-68, 1996).

25

EXENDIN

Exendins are peptides that are found in the venom of the Gila-monster, a lizard endogenous to Arizona, and the Mexican Beaded Lizard. Exendin-3 is present in the venom of *Heloderma horridum*, and exendin-4 is present in the venom of *Heloderma*

30 *suspectum* (Eng, J., et al., J. Biol. Chem., 265:20259-62, 1990; Eng, J., et al., J. Biol. Chem., 267:7402-05, 1992). The exendins have some sequence similarity to several

members of the glucagon-like peptide family, with the highest homology, 53%, being to GLP-1 (Goke, et al., J. Biol. Chem., 268:19650-55, 1993).

Exendin-4 is a potent agonist at GLP-1 receptors on insulin-secreting TC1 cells, at dispersed acinar cells from guinea pig pancreas, and at parietal cells from stomach; the peptide also stimulates somatostatin release and inhibits gastrin release in isolated stomachs (Goke, et al., J. Biol. Chem. 268:19650-55, 1993; Schepp, et al., Eur. J. Pharmacol., 69:183-91, 1994; Eissele, et al., Life Sci., 55:629-34, 1994). Exendin-3 and exendin-4 were found to be GLP-1 agonists in stimulating cAMP production in, and amylase release from, pancreatic acinar cells (Malhotra, R., et al., Regulatory Peptides, 41:149-56, 1992; Raufman, et al., J. Biol. Chem. 267:21432-37, 1992; Singh, et al., Regul. Pept. 53:47-59, 1994). The use of the insulinotropic activities of exendin-3 and exendin-4 for the treatment of diabetes mellitus and the prevention of hyperglycemia has been proposed (Eng, U.S. Patent No. 5,424,286).

Truncated exendin peptides such as exendin[9-39], a carboxyamidated molecule, and fragments 3-39 through 9-39 have been reported to be potent and selective antagonists of GLP-1 (Goke, et al., J. Biol. Chem., 268:19650-55, 1993; Raufman, J.P., et al., J. Biol. Chem. 266:2897-902, 1991; Schepp, W., et al., Eur. J. Pharm. 269:183-91, 1994; Montrose-Rafizadeh, et al., Diabetes, 45(Suppl. 2):152A, 1996). Exendin[9-39] blocks endogenous GLP-1 in vivo, resulting in reduced insulin secretion. Wang, et al., J. Clin. Invest., 95:417-21, 1995; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996). The receptor apparently responsible for the insulinotropic effect of GLP-1 has been cloned from rat pancreatic islet cells (Thorens, B., Proc. Natl. Acad. Sci. USA 89:8641-8645, 1992). Exendins and exendin[9-39] bind to the cloned GLP-1 receptor (rat pancreatic -cell GLP-1 receptor: Fehmann HC, et al., Peptides 15 (3): 453-6, 1994; human GLP-1 receptor: Thorens B, et al., Diabetes 42 (11): 1678-82, 1993). In cells transfected with the cloned GLP-1 receptor, exendin-4 is an agonist, i.e., it increases cAMP, while exendin[9-39] is an antagonist, i.e., it blocks the stimulatory actions of exendin-4 and GLP-1. Id.

Exendin[9-39] also acts as an antagonist of the full length exendins, inhibiting stimulation of pancreatic acinar cells by exendin-3 and exendin-4 (Raufman, et al., J. Biol. Chem. 266:2897-902, 1991; Raufman, et al., J. Biol. Chem., 266:21432-37, 1992). Exendin[9-39] inhibits the stimulation of plasma insulin levels by exendin-4, and inhibits

the somatostatin release-stimulating and gastrin release-inhibiting activities of exendin-4 and GLP-1 (Kolligs, F., et al., Diabetes, 44:16-19, 1995; Eissele, et al., Life Sciences, 55:629-34, 1994). Exendin-4, administered through the jugular vein, has been reported to induce an increase in systolic, diastolic and mean arterial blood pressure, and in heart rate
5 (Barragán, et al., Regul. Pep. 67:63-68, 1996).

Exendins have recently been found to inhibit gastric emptying (U.S. Patent Application Serial No. 08/694,954, filed August 8, 1996, which enjoys common ownership with the present invention and is hereby incorporated by reference). Exendin[9-39] has been used to investigate the physiological relevance of central GLP-1
10 in control of food intake (Turton, M.D. et al., Nature, 379:69-72, 1996). GLP-1 administered by intracerebroventricular (ICV) injection inhibits food intake in rats. This satiety-inducing effect of GLP-1 delivered by intracerebroventricular injection is reported to be inhibited by ICV injection of exendin[9-39] (Turton, supra). However, it has been reported that GLP-1 does not inhibit food intake in mice when administered by peripheral
15 injection (Turton, M.D., Nature 379:69-72, 1996; Bhavsar, S.P., Soc. Neurosci. Abstr. 21:460 (188.8), 1995). Administration of exendins and exendin agonists has also recently been found to reduce food intake (U.S. Provisional Patent Application Serial No. 60/034,905, filed January 7, 1997, which enjoys common ownership with the present invention and is hereby incorporated by reference).

20 DIURETICS

Agents that increase urine flow, or diuretics, are useful for treating conditions or disorders that are associated with toxic hypervolemic states. Such conditions or disorders include renal failure, congestive heart failure, nephrotic syndrome, cirrhosis, pulmonary
25 edema, and hypertension. Diuretics are also employed to treat conditions in pregnancy, such as pre-eclampsia and eclampsia. Further uses of diuretics include their use to reduce volume before some surgical procedures such as ocular surgery and neurosurgery.

One difficulty encountered with many diuretics such as thiazides, loop diuretics, carbonic anhydrase inhibitors, and osmotic diuretics, is that although they may be
30 employed to increase sodium excretion, they also result in an increase of urinary potassium loss. Examples of the effects of potassium loss include muscular weakness,

paralysis (including the paralysis of respiratory muscles), electrocardiographic abnormalities, cardiac dysrhythmia, and cardiac arrest.

Another difficulty encountered with some diuretics is their slow rate of action, which is not conducive to their use in an emergency setting.

5 Thus, there is a need for a method of increasing urine flow that does not deplete potassium concentration in the patient and which has a rapid mode of action. Such methods, and compounds and compositions which are useful therefore, have been invented and are described and claimed herein.

Inotropic Compounds

10 Compounds that induce inotropic effects (e.g., increase of force of contraction of the heart) have been recognized as being useful for the treatment of, for example, congestive heart failure. Congestive heart failure, which is one of the most common causes of death and disability in industrialized nations, has a mortality rate of about 50% at five years (Goodman and Gilman's The Pharmacological Basis of Therapeutics, 9th Ed.
15 McGraw Hill, New York, pp. 809-838). Inotropic agents currently in clinical use include digitalis, sympathomimetic amines and amrinone (Harrison's Principles of Internal Medicine, 12th Edition, 1991, McGraw Hill, New York, pp. 894-899).

 Digoxin, a cardiac glycoside, an ancient but effective therapy for cardiac failure, was initially derived from the foxglove leaf, *Digitalis purpurea* and *Digitalis lanata*.
20 Cardiac glycosides are potent and highly selective inhibitors of the active transport of sodium and potassium ions across cell membranes (Goodman and Gilman, supra). Cardiac glycosides have been reported to increase the velocity of shortening of cardiac muscle, resulting in an improvement in ventricular function; this effect has been reported to be due to an increase in the availability during systole of cytosolic Ca^{2+} to interact with
25 contractile proteins to increase the velocity and extent of sarcomere shortening (Goodman and Gilman, supra).

 Digoxin and related cardiac glycosides (e.g., digoxin) have useful durations of action because their excretion, mainly via the kidneys, results in plasma $t_{1/2}$ of 1.5-5 days. But the therapeutic index of these drugs is very low with mildly toxic:minimally-effective
30 dose ratio being 2:1 and lethal:minimally-effective dose ratio being between 5:1 and 10:1. Urinary potassium loss due to use of thiazide and loop diuretics may seriously enhance

the dangers of digitalis intoxication, including susceptibility to cardiac arrhythmia, and potassium-sparing diuretics are often necessary. Slow elimination of cardiac glycosides can prolong the period of jeopardy during digitalis intoxication, which has been reported to occur in 20% of hospital patients on these drugs. Absorption and onset of action for all
5 cardiac glycosides except ouabain is somewhat prolonged, and this may be a disadvantage in emergency cardiac conditions.

Sympathomimetic amines, which generally include epinephrine, isoproterenol, dopamine and dobutamine, can be useful in an acute setting to stimulate myocardial contractility, but they usually require constant intravenous infusion and continuous
10 intensive monitoring of the patient. They typically lose their effectiveness after 8 hours, apparently due to receptor downregulation.

Amrinone, a noncatecholamine, non-glycoside agent also requires continuous intravenous administration.

This description of available inotropic agents illustrates the need for, and
15 desirability of, therapies that are (1) inotropic, with (2) rapid onset of action, with (3) prolonged duration of action (including a persistent effect, with absence of tachyphylaxis), with (4) low toxicity (a high ratio of toxic to therapeutic dose), with (5) rapid and profound diuretic effect, with (6) a sparing of urinary potassium loss, and with (7) a convenient (non-intravenous) route of administration. We have discovered that exendin and GLP-1
20 fulfill these criteria.

SUMMARY OF THE INVENTION

The present invention concerns the surprising discovery that exendins, GLP-1, and agonists of these compounds have rapid inotropic and diuretic effects. Although GLP-1
25 has been reported to not have a diuretic effect when administered peripherally, we have found, surprisingly, that GLP-1 does in fact have a diuretic effect after peripheral administration. This diuretic effect of exendins, GLP-1, and exendin and GLP-1 agonists, is accompanied by an increase in urinary sodium concentration. This diuretic effect is also accompanied by a decrease in urinary potassium concentration which is unanticipated as
30 many diuretics have been found to cause a profound increase in urinary potassium concentration.

The present invention is directed to novel methods for increasing urine flow comprising the administration of an exendin, for example, exendin-3 [SEQ ID NO. 1: His Ser Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH₂], or exendin-4
5 [SEQ ID NO. 2: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH₂], or other compounds which effectively bind to the receptor at which exendin exerts its action on increasing urine flow (exendin agonists). The present invention is also directed to novel methods for increasing urine flow comprising the administration of GLP-
10 1 [SEQ ID NO. 3: His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg-NH₂], or other compounds which effectively bind to the receptor at which GLP-1 exerts its action on increasing urine flow (GLP-1 agonists).

In a first aspect, the invention features a method of increasing urine flow in an
15 individual comprising administering to the individual a therapeutically effective amount of an exendin or an exendin agonist. In one preferred aspect, said exendin is exendin-3. More preferably, said exendin is exendin-4. By an exendin agonist is meant a compound that mimics the effects of exendin on increasing urine flow, increasing sodium excretion, and/or decreasing urinary potassium concentration, (the potassium concentration in
20 excreted urine) by binding to the receptor or receptors where exendin causes this effect. Certain novel exendin agonist compounds are described in U.S. Provisional Patent Application Serial No. 60/055,404, filed August 8, 1997, which enjoys common ownership with the present invention and is hereby incorporated by this reference. Certain other novel exendin agonist compounds are described in U.S. Provisional Patent
25 Application Serial Nos. 60/066,029 and 60/065,442, both filed November 14, 1997 which enjoy common ownership with the present invention and are hereby incorporated by this reference. Preferred exendin agonist compounds include those described in U.S. Provisional Patent Application Serial Nos. 60/055,404 and 60/065,442.

In one preferred aspect the exendin or exendin agonist used in the methods of the
30 present invention is exendin-4. In another preferred aspect, the exendin is exendin-3. In other preferred aspects, the exendin or exendin agonist is a compound of the formula (I)

[SEQ ID NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

- 5 Xaa₁ is His, Arg or Tyr;
Xaa₂ is Ser, Gly, Ala or Thr;
Xaa₃ is Asp or Glu;
Xaa₅ is Ala or Thr;
Xaa₆ is Ala, Phe, Tyr or naphthylalanine;
- 10 Xaa₇ is Thr or Ser;
Xaa₈ is Ala, Ser or Thr;
Xaa₉ is Asp or Glu;
Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
Xaa₁₁ is Ala or Ser;
- 15 Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
Xaa₁₅ is Ala or Glu;
Xaa₁₆ is Ala or Glu;
- 20 Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
Xaa₂₁ is Ala or Leu;
Xaa₂₂ is Phe, Tyr or naphthylalanine;
- 25 Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
or Met;
Xaa₂₄ is Ala, Glu or Asp;
Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
Xaa₂₆ is Ala or Leu;
- 30 Xaa₂₇ is Ala or Lys;
Xaa₂₈ is Ala or Asn;

Z₁ is -OH,

-NH₂,

Gly-Z₂,

Gly Gly-Z₂,

5 Gly Gly Xaa₃₁-Z₂,

Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

10 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂, or

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈ Xaa₃₉-Z₂;

wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently

15 selected from the group consisting of Pro,

homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine and

N-alkylalanine; Xaa₃₉ is Ser, Thr or Tyr; and

Z₂ is -OH or -NH₂; and pharmaceutically acceptable salts thereof;

20 provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃,

Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇, and Xaa₂₈ are

Ala; and provided also that the compound is not exendin-3 [SEQ ID NO. 1] or exendin-4

[SEQ ID NO. 2]. In other aspects of the invention, the increase in urine flow is

accompanied by an increase in sodium excretion in said individual. In most preferred

25 aspects, the increase in urine flow does not increase urinary potassium concentration in said individual.

In other embodiments of the invention, a method is provided for decreasing the concentration of potassium in the urine of an individual comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

30 In yet another aspect of the invention, a method is provided for preventing or alleviating a condition or disorder associated with toxic hypervolemia in an individual,

comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

By condition or disorder associated with toxic hypervolemia is meant any condition or disorder in a subject that is either caused by, complicated by, or aggravated by a relatively high extracellular volume. Such conditions or disorders include, but are not limited to, renal failure, congestive heart failure, nephrotic syndrome, pulmonary edema, cirrhosis, and hypertension.

The present invention also provides a method of inducing rapid diuresis in an individual comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist. One preferred use of this method is in preparation of a patient for a surgical procedure where a reduction in extracellular volume is desired, such as in some ocular surgical procedures or in some neurosurgical procedures. Thus, the present invention provides a method of preparing an individual for a surgical procedure comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist. Preferably, said exendin or exendin agonist is administered to said individual before said surgical procedure.

In other preferred aspects, a method is provided for increasing renal plasma flow and glomerular filtration rate in an individual comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

In yet other preferred aspects, a method is provided for treating pre-eclampsia or eclampsia of pregnancy in an individual comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

The preferred mode of administration of said exendin or exendin agonist is by peripheral (subcutaneous or intravenous) administration. Preferably, said exendin or exendin agonist is administered subcutaneously. Preferably, about 1 g - 30 g to about 10-20 mg of the exendin or exendin agonist is administered per dose. More preferably, about 30 g to about 10 mg, or about 300 g to about 5mg of the exendin or exendin agonist is administered per dose. Most preferably, about 30 g to about 1 mg of the exendin or exendin agonist is administered per dose.

In other preferred aspects, said peripheral administration is selected from the group consisting of buccal, nasal, pulmonary, oral, intraocular, rectal, and transdermal

administration.

The present invention also provides pharmaceutical compositions for use in the treatment of conditions or disorders associated with hypervolemia comprising a therapeutically effective amount of an exendin or exendin agonist in association with a
5 pharmaceutically acceptable carrier.

In yet other aspects, the invention provides pharmaceutical compositions for use in increasing urine flow in an individual comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

In further aspects, the invention provides pharmaceutical compositions for use in
10 treating pre-eclampsia or eclampsia of pregnancy in an individual comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

Preferably, these pharmaceutical compositions comprise exendin-3. More preferably, these pharmaceutical compositions comprise exendin-4.

15 Preferably, these pharmaceutical compositions comprise an exendin agonist of formula I [SEQ ID NO. 4].

The present invention is also directed to novel methods for increasing urine flow comprising the administration of GLP-1.

In one embodiment the invention features a method of increasing urine flow in an
20 individual comprising administering to the individual a therapeutically effective amount of GLP-1 or GLP-1 agonist. By GLP-1 agonist is meant a compound that mimics the effects of GLP-1 on increasing urine flow, increasing sodium excretion, and/or decreasing urinary potassium concentration, by binding to the receptor or receptors where GLP-1 causes this effect. Certain GLP-1 agonists are described in Chen et al., U.S. Patent No.
25 5,512,549, issued April 30, 1996, entitled Glucagon-Like Insulinotropic Peptide Analogs, Compositions and Methods of Use. Other GLP-1 agonists are described in Johnson et al., U.S. Patent No. 5,574,008, issued November 12, 1996, entitled, Biologically Active Fragments of Glucagon-Like Insulinotropic Peptide. Still other GLP-1 agonists are described in Buckley et al., U.S. Patent No. 5,545,618, issued August 13, 1996, entitled
30 GLP-1 Analogs Useful for Diabetes Treatment. All three referenced U.S. patents are incorporated herein by this reference.

In other aspects of the invention, the increase in urine flow is accompanied by an increase in sodium excretion in said individual. In most preferred aspects, the increase in urine flow does not increase urinary potassium concentration in said individual.

In other embodiments of the invention, a method is provided for decreasing the
5 concentration of potassium in the urine of an individual comprising administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist.

In yet another aspect of the invention, a method is provided for preventing or alleviating a condition or disorder associated with toxic hypervolemia in an individual, comprising administering to said individual a therapeutically effective amount of GLP-1
10 or a GLP-1 agonist.

The present invention also provides a method of inducing rapid diuresis in an individual comprising administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist. One preferred use of this method is in preparation of patient for surgical procedures where a reduction in extracellular volume is desired, such as in
15 some ocular surgical procedures and some neurosurgical procedures. Thus, the present invention provides a method of preparing an individual for a surgical procedure comprising administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist. Preferably, said GLP-1 or GLP-1 agonist is administered to said individual before said surgical procedure.

20 In other preferred aspects, a method is provided for increasing renal plasma flow and glomerular filtration rate in an individual comprising administering to said individual a therapeutically effective amount of GLP-1 or GLP-1 agonist.

In yet other preferred aspects, a method is provided for treating pre-eclampsia or eclampsia of pregnancy in an individual comprising administering to said individual a
25 therapeutically effective amount of GLP-1 or GLP-1 agonist.

The preferred mode of administration of said GLP-1 or GLP-1 agonist is by peripheral administration. Preferably, said GLP-1 or GLP-1 agonist is administered subcutaneously or intravenously. Preferably, about 1 g-30 g to about 10-20 mg of GLP-1 or GLP-1 agonist is administered per dose. More preferably, about 30 g to about 10mg,
30 or about 300 g to about 5mg of GLP-1 or GLP-1 agonist is administered per dose. Most preferably, about 30 g to about 1 mg of GLP-1 or GLP-1 agonist is administered per dose.

In other preferred aspects, said peripheral administration is selected from the group consisting of buccal, nasal, pulmonary, oral, intraocular, rectal, and transdermal administration.

5 The present invention also provides pharmaceutical compositions for use in the treatment of conditions or disorders associated with hypervolemia comprising a therapeutically effective amount of GLP-1 or a GLP-1 agonist in association with a pharmaceutically acceptable carrier.

10 In yet other aspects, the invention provides pharmaceutical compositions for use in increasing urine flow in an individual comprising a therapeutically effective amount of GLP-1 or a GLP-1 agonist in association with a pharmaceutically acceptable carrier.

15 In further aspects, the invention provides pharmaceutical compositions for use in treating pre-eclampsia or eclampsia of pregnancy in an individual comprising a therapeutically effective amount of GLP-1 or a GLP-1 agonist in association with a pharmaceutically acceptable carrier.

20 The present invention also features methods for inducing an inotropic effect in an individual comprising administering a therapeutically effective amount of an exendin or an exendin agonist, or GLP 1 or a GLP 1 agonist. Thus, in one aspect, is provided a method for increasing cardiac contractility in an individual comprising administering a therapeutically effective amount of an exendin, an exendin agonist, GLP-1 or a GLP-1 agonist.

25 In a related aspect, a method is provided for treating a condition or disorder that can be alleviated by increasing cardiac contractility in an individual comprising administering a therapeutically effective amount of an exendin, an exendin agonist, GLP-1 or a GLP-1 agonist. Such conditions or disorders include congestive heart failure, pulmonary and systemic edema, and renal failure. Preferably, said condition or disorder is congestive heart failure.

Preferably, said exendin is to be used in those methods exendin-3. More preferably, said exendin is exendin-4.

30 Preferably, the exendin agonist to be used in those methods is an exendin agonist of formula (I) [SEQ ID NO. 4].

In preferred aspects, said exendin, exendin agonist, GLP-1, or a GLP-1 agonist to be used in these methods is administered peripherally using the doses described herein.

Preferably, said peripheral administration is selected from the group consisting of buccal, nasal, pulmonary, oral, intraocular, rectal, and transdermal administration.

5 In another preferred aspect, said exendin, exendin agonist, GLP-1, or a GLP-1 agonist is administered subcutaneously or intravenously.

Also provided in the present invention are pharmaceutical compositions for use in the treatment of a condition or disorder that can be alleviated by increasing cardiac contractility comprising a therapeutically effective amount of an exendin, an exendin
10 agonist, GLP-1 or a GLP-1 agonist in association with a pharmaceutically acceptable carrier. Preferably, said exendin is exendin-3. More preferably, said exendin is exendin-4. Preferably, these pharmaceutical compositions comprise an exendin agonist of formula I [SEQ ID NO. 4].

15 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1(A-B) is a graphical depiction of the response of mean arterial pressure (MAP) to GLP-1. (A) MAP is presented as % of predose values measured over the 30 minutes prior to drug administration; (B) Dose-response curve for effects of GLP-1 of
20 MAP. The response plotted is the incremental area under the curve from 0 to 2 hours after the bolus dose.

Figure 2 is a graphical depiction of the inotropic response to GLP-1. The rate of change of blood pressure (dp/dt) is indicative of cardiac contractility, which increased in response to a subcutaneous injection of GLP-1 given to conscious rats.

Figure 3(A-B) is a graphical depiction of the response of urine flow to intravenous
25 bolus doses of GLP-1. (A) Urine flow was measured at 15 minute intervals and presented as % of predose values measured over the 30 minutes prior to drug administration; (B) Dose-response curve for effects of GLP-1 on urine flow. The response plotted is the percent change in flow from 0 to 15 minutes after the bolus dose relative to the flow over the previous 30 minutes.

30 Figure 4(A-B) is a graphical depiction of the response of sodium excretion to intravenous bolus doses of GLP-1. (A) Sodium excretion was measured at 15 minute

intervals and presented as % of predose values measured over the 30 minutes prior to drug administration; (B) Dose-response curve for effects of GLP-1 on sodium excretion. The response plotted is the percent change in sodium excretion from 0 to 15 minutes after the bolus dose relative to excretion over the previous 30 minutes.

5 Figure 5(A-B) is a graphical depiction of the response of urinary potassium concentration to intravenous bolus doses of GLP-1. (A) Urinary potassium concentration was measured at 15 minute intervals and presented as % of predose values measured over the 30 minutes prior to drug administration; (B) Dose-response curve for effects of GLP-1 on urinary potassium concentration. The response plotted is the percent change in urinary
10 potassium concentration from 0 to 15 minutes after the bolus dose relative to the urinary potassium concentration over the previous 30 minutes.

 Figure 6(A-B) is a graphical depiction of the response of mean arterial pressure (MAP) to exendin 4. (A) MAP is presented as % of predose values measured over the 30 minutes prior to drug administration; (B) Dose-response curve for effects of exendin on
15 MAP. The response plotted is the incremental area under the curve from 0 to 2 hours after the bolus dose.

 Figure 7 is a graphical depiction of the inotropic response to exendin-4. The rate of change of blood pressure (dP/dt) is indicative of cardiac contractility, which increased in response to a subcutaneous injection of exendin-4 given to conscious rats.

20 Figure 8(A-B) is a graphical depiction of the response of urine flow to intravenous bolus doses of exendin-4. (A) Urine flow was measured at 15 minute intervals; (B) Dose-response curve for effects of exendin-4 on urine flow. The response plotted is urine flow from 0 to 15 minutes after the bolus dose.

 Figure 9(A-B) is a graphical depiction of the response of sodium excretion to
25 intravenous bolus doses of exendin-4. (A) Sodium excretion was measured at 15 minute intervals and presented as % of predose values measured over the 30 minutes prior to drug administration; (B) Dose-response curve for effects of exendin-4 on sodium excretion. The response plotted is the percent change in sodium excretion from 0 to 15 minutes after the bolus dose relative to excretion over the previous 30 minutes.

30 Figure 10(A-B) is a graphical depiction of the response of urinary potassium concentration to intravenous bolus doses of exendin-4. (A) Urinary potassium

concentration was measured at 15 minute intervals and presented as % of predose values measured over the 30 minutes prior to exendin-4 administration; (B) Dose-response curve for effects of exendin on urinary potassium concentration. The response plotted is the incremental area under the curve from 0 to 2 hours after the bolus dose.

5


DETAILED DESCRIPTION OF THE INVENTION

The extendins, GLP-1, and analogs and agonists thereof of this invention are useful in view of their pharmacological properties. Activity as exendin or GLP-1 analogs or agonists can be indicated by activity in the assays described below. Effects of extendins or GLP-1 agonists thereof on reducing food intake can be identified, evaluated, or screened for, using the methods described in the Examples below, or other methods known in the art for determining effects on urine flow, or sodium or potassium excretion.

Although exendin-4 was found to have a hypertensive effect, when administered in conjunction with an agent that regulated blood pressure, the diuretic effect was still evident, indicating a diuretic effect of exendin-4 that was not entirely attributable to its hypertensive effect.

Exendin Agonist Compounds

Exendin agonist compounds include those described in U.S. Provisional Patent Application Nos. 60/055,404; 60/066,029; and 60/065,442. Preferred exendin agonist compounds include peptide compounds of the formula (I) [SEQ ID NO. 4]:



25

Xaa₁ is His, Arg or Tyr;

Xaa₂ is Ser, Gly, Ala or Thr;

Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

30 Xaa₆ is Ala, Phe, Tyr or naphthylalanine;

Xaa₇ is Thr or Ser;

- Xaa₈ is Ala, Ser or Thr;
 Xaa₉ is Asp or Glu;
 Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
 Xaa₁₁ is Ala or Ser;
 5 Xaa₁₂ is Ala or Lys;
 Xaa₁₃ is Ala or Gln;
 Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
 Xaa₁₅ is Ala or Glu;
 Xaa₁₆ is Ala or Glu;
 10 Xaa₁₇ is Ala or Glu;
 Xaa₁₉ is Ala or Val;
 Xaa₂₀ is Ala or Arg;
 Xaa₂₁ is Ala or Leu;
 Xaa₂₂ is Phe, Tyr or naphthylalanine;
 15 Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
 or Met;
 Xaa₂₄ is Ala, Glu or Asp;
 Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
 Xaa₂₆ is Ala or Leu;
 20 Xaa₂₇ is Ala or Lys;
 Xaa₂₈ is Ala or Asn;
 Z₁ is -OH,
 -NH₂,
 Gly-Z₂,
 25 Gly Gly-Z₂,
 Gly Gly Xaa₃₁-Z₂,
 Gly Gly Xaa₃₁ Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
 30 Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂, or
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈ Xaa₃₉-Z₂;

wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently

5 selected from the group consisting of Pro,
homoproline, 3Hyp, 4Hyp, thioproline,
N-alkylglycine, N-alkylpentylglycine and
N-alkylalanine; Xaa₃₉ is Ser, Thr or Tyr; and
Z₂ is -OH or -NH₂; and pharmaceutically acceptable salts thereof;

10 provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃,
Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇, and Xaa₂₈ are
Ala; and provided also that the compound is not exendin-3 or exendin-4.

Preferred N-alkyl groups for N-alkylglycine, N-alkylpentylglycine and N-
alkylalanine include lower alkyl groups preferably of 1 to about 6 carbon atoms, more
15 preferably of 1 to 4 carbon atoms. Suitable compounds include those identified in
Examples 4-64 [SEQ ID NOS. 5 to 65], as well as those compounds identified in
Examples 65 and 66.

Preferred exendin agonist compounds include those wherein Xaa₁ is His or Tyr.
More preferably Xaa₁ is His.

20 Preferred are those compounds wherein Xaa₂ is Gly.

Preferred are those compounds wherein Xaa₁₄ is Leu, pentylglycine or Met.

Preferred compounds are those wherein Xaa₂₅ is Trp or Phe.

Preferred compounds are those where Xaa₆ is Phe or naphthylalanine; Xaa₂₂ is Phe
of naphthylalanine and

25 Xaa₂₃ is Ile or Val.

Preferred are compounds wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently
selected from Pro, homoproline, thioproline and N-alkylalanine.

Preferably Z₁ is -NH₂.

Preferably Z₂ is -NH₂.

30 According to one aspect, preferred are compounds of formula (I) wherein Xaa₁ is
His or Tyr, more preferably His; Xaa₂ is Gly; Xaa₆ is Phe or naphthylalanine; Xaa₁₄ is Leu,

pentylglycine or Met; Xaa₂₂ is Phe or naphthylalanine; Xaa₂₃ is Ile or Val; Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from Pro, homoproline, thioproline or N-alkylalanine. More preferably Z₁ is -NH₂.

According to an especially preferred aspect, especially preferred compounds
 5 include those of formula (I) wherein: Xaa₁ is His or Arg; Xaa₂ is Gly or Ala; Xaa₃ is Asp or Glu; Xaa₄ is Ala or Thr; Xaa₅ is Ala, Phe or naphthylalanine; Xaa₆ is Thr or Ser; Xaa₈ is Ala, Ser or Thr; Xaa₉ is Asp or Glu; Xaa₁₀ is Ala, Leu or pentylglycine; Xaa₁₁ is Ala or Ser; Xaa₁₂ is Ala or Lys; Xaa₁₃ is Ala or Gln; Xaa₁₄ is Ala, Leu or pentylglycine; Xaa₁₅ is Ala or Glu; Xaa₁₆ is Ala or Glu; Xaa₁₇ is Ala or Glu; Xaa₁₉ is Ala or Val; Xaa₂₀ is Ala or Arg; Xaa₂₁ is Ala or Leu; Xaa₂₂ is Phe or naphthylalanine; Xaa₂₃ is Ile, Val or tert-
 10 butylglycine; Xaa₂₄ is Ala, Glu or Asp; Xaa₂₅ is Ala, Trp or Phe; Xaa₂₆ is Ala or Leu; Xaa₂₇ is Ala or Lys; Xaa₂₈ is Ala or Asn; Z₁ is -OH, -NH₂, Gly-Z₂, Gly Gly-Z₂, Gly Gly Xaa₃₁-Z₂, Gly Gly Xaa₃₁ Ser-Z₂, Gly Gly Xaa₃₁ Ser Ser-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂; Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ being independently Pro homoproline, thioproline or N-methylalanine; and Z₂ being -OH or -NH₂; provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala. Especially preferred compounds include those having the
 15 amino acid sequence of SEQ ID NOS. 6-27.

According to an especially preferred aspect, provided are compounds where Xaa₁₄ is Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and Xaa₂₅ is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will be less susceptible to oxidative degradation, both *in vitro* and *in vivo*, as well as during
 25 synthesis of the compound.

Definitions


In accordance with the present invention and as used herein, the following terms are defined to have the following meanings, unless explicitly stated otherwise.

30 The term "amino acid" refers to natural amino acids, unnatural amino acids, and amino acid analogs, all in their D and L stereoisomers if their structure allow such

stereoisomeric forms. Natural amino acids include alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp), cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), Lysine (Lys), methionine (Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr),
 5 typtophan (Trp), tyrosine (Tyr) and valine (Val). Unnatural amino acids include, but are not limited to azetidinecarboxylic acid, 2-aminoadipic acid, 3-aminoadipic acid, beta-alanine, aminopropionic acid, 2-aminobutyric acid, 4-aminobutyric acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-aminoisobutyric acid, 2-aminopimelic acid, tertiary-butylglycine, 2,4-diaminoisobutyric acid, desmosine, 2,2'-
 10 diaminopimelic acid, 2,3-diaminopropionic acid, N-ethylglycine, N-ethylasparagine, homoproline, hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-hydroxyproline, isodesmosine, allo-isoleucine, N-methylalanine, N-methylglycine, N-methylisoleucine, N-methylpentylglycine, N-methylvaline, naphthalanine, norvaline, norleucine, ornithine, pentylglycine, pipercolic acid and thioproline. Amino acid analogs include the natural and
 15 unnatural amino acids which are chemically blocked, reversibly or irreversibly, or modified on their N-terminal amino group or their side-chain groups, as for example, methionine sulfoxide, methionine sulfone, S-(carboxymethyl)-cysteine, S-(carboxymethyl)-cysteine sulfoxide and S-(carboxymethyl)-cysteine sulfone.

The term "amino acid analog" refers to an amino acid wherein either the C-
 20 terminal carboxy group, the N-terminal amino group or side-chain functional group has been chemically codified to another functional group. For example, aspartic acid-(beta-methyl ester) is an amino acid analog of aspartic acid; N-ethylglycine is an amino acid analog of glycine; or alanine carboxamide is an amino acid analog of alanine.

The term "amino acid residue" refers to radicals having the structure: (1) -C(O)-R-
 25 NH-, wherein R typically is -CH(R)-, wherein R is an amino acid side chain, typically H or a carbon containing substituent; or (2)



wherein p is 1, 2 or 3 representing the azetidinecarboxylic acid, proline or pipecolic acid residues, respectively.

The term "lower" referred to herein in connection with organic radicals such as alkyl groups defines such groups with up to and including about 6, preferably up to and including 4 and advantageously one or two carbon atoms. Such groups may be straight chain or branched chain.

"Pharmaceutically acceptable salt" includes salts of the compounds described herein derived from the combination of such compounds and an organic or inorganic acid. In practice the use of the salt form amounts to use of the base form. The compounds are useful in both free base and salt form.

In addition, the following abbreviations stand for the following:

"ACN" or "CH₃CN" refers to acetonitrile.

"Boc", "tBoc" or "Tboc" refers to t-butoxy carbonyl.

"DCC" refers to N,N'-dicyclohexylcarbodiimide.

"Fmoc" refers to fluorenylmethoxycarbonyl.

"HBTU" refers to 2-(1H-benzotriazol-1-yl)-

1,1,3,3-tetramethyluronium hexafluorophosphate.

"HOBT" refers to 1-hydroxybenzotriazole monohydrate.

"homoP" or hPro" refers to homoproline.

"MeAla" or "Nme" refers to N-methylalanine.

"naph" refers to naphthylalanine.

"pG" or pGly" refers to pentylglycine.

"tBuG" refers to tertiary-butylglycine.

"ThioP" or tPro" refers to thioproline.

3Hyp" refers to 3-hydroxyproline

4Hyp" refers to 4-hydroxyproline

NAG" refers to N-alkylglycine

NAPG" refers to N-alkylpentylglycine

"Norval" refers to norvaline

"Norleu" refers to norleucine

Preparation of Compounds

Compounds such as the exendins and exendin agonists described herein may be prepared using standard solid-phase peptide synthesis techniques and preferably an automated or semiautomated peptide synthesizer. Typically, using such techniques, an -N-carbamoyl protected amino acid and an amino acid attached to the growing peptide chain on a resin are coupled at room temperature in an inert solvent such as dimethylformamide, N-methylpyrrolidinone or methylene chloride in the presence of coupling agents such as dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in the presence of a base such as diisopropylethylamine. The -N-carbamoyl protecting group is removed from the resulting peptide-resin using a reagent such as trifluoroacetic acid or piperidine, and the coupling reaction repeated with the next desired N-protected amino acid to be added to the peptide chain. Suitable N-protecting groups are well known in the art, with t-butyloxycarbonyl (tBoc) and fluorenylmethoxycarbonyl (Fmoc) being preferred herein.

The solvents, amino acid derivatives and 4-methylbenzhydryl-amine resin used in the peptide synthesizer may be purchased from Applied Biosystems Inc. (Foster City, CA). The following side-chain protected amino acids may be purchased from Applied Biosystems, Inc.: Boc-Arg(Mts), Fmoc-Arg(Pmc), Boc-Thr(Bzl), Fmoc-Thr(t-Bu), Boc-Ser(Bzl), Fmoc-Ser(t-Bu), Boc-Tyr(BrZ), Fmoc-Tyr(t-Bu), Boc-Lys(Cl-Z), Fmoc-Lys(Boc), Boc-Glu(Bzl), Fmoc-Glu(t-Bu), Fmoc-His(Trt), Fmoc-Asn(Trt), and Fmoc-Gln(Trt). Boc-His(BOM) may be purchased from Applied Biosystems, Inc. or Bachem Inc. (Torrance, CA). Anisole, dimethylsulfide, phenol, ethanedithiol, and thioanisole may be obtained from Aldrich Chemical Company (Milwaukee, WI). Air Products and Chemicals (Allentown, PA) supplies HF. Ethyl ether, acetic acid and methanol may be purchased from Fisher Scientific (Pittsburgh, PA).

Solid phase peptide synthesis may be carried out with an automatic peptide synthesizer (Model 430A, Applied Biosystems Inc., Foster City, CA) using the NMP/HOBt (Option 1) system and tBoc or Fmoc chemistry (see, Applied Biosystems User's Manual for the ABI 430A Peptide Synthesizer, Version 1.3B July 1, 1988, section 6, pp. 49-70, Applied Biosystems, Inc., Foster City, CA) with capping. Boc-peptide-resins may be cleaved with HF (-5°C to 0°C, 1 hour). The peptide may be extracted from the resin with alternating water and acetic acid, and the filtrates lyophilized. The Fmoc-

peptide resins may be cleaved according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc., 1990, pp. 6-12). Peptides may be also be assembled using an Advanced Chem Tech Synthesizer (Model MPS 350, Louisville, Kentucky).

5 Peptides may be purified by RP-HPLC (preparative and analytical) using a Waters Delta Prep 3000 system. A C4, C8 or C18 preparative column (10 μ , 2.2 x 25 cm; Vydac, Hesperia, CA) may be used to isolate peptides, and purity may be determined using a C4, C8 or C18 analytical column (5 μ , 0.46 x 25 cm; Vydac). Solvents (A=0.1% TFA/water and B=0.1% TFA/CH₃CN) may be delivered to the analytical column at a flowrate of 1.0
10 ml/min and to the preparative column at 15 ml/min. Amino acid analyses may be performed on the Waters Pico Tag system and processed using the Maxima program. Peptides may be hydrolyzed by vapor-phase acid hydrolysis (115°C, 20-24 h). Hydrolysates may be derivatized and analyzed by standard methods (Cohen, et al., The Pico Tag Method: A Manual of Advanced Techniques for Amino Acid Analysis, pp.
15 11-52, Millipore Corporation, Milford, MA (1989)). Fast atom bombardment analysis may be carried out by M-Scan, Incorporated (West Chester, PA). Mass calibration may be performed using cesium iodide or cesium iodide/glycerol. Plasma desorption ionization analysis using time of flight detection may be carried out on an Applied Biosystems Bio-Ion 20 mass spectrometer. Electrospray mass spectroscopy may be carried out on a VG-
20 Trio machine.

Peptide compounds useful in the invention may also be prepared using recombinant DNA techniques, using methods now known in the art. See, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d Ed., Cold Spring Harbor (1989). Non-peptide compounds useful in the present invention may be prepared by art-known
25 methods. For example, phosphate-containing amino acids and peptides containing such amino acids, may be prepared using methods known in the art. See, e.g., Bartlett and Landen, Biorg. Chem. 14:356-377 (1986).

Exendin or GLP-1 agonist analogs or derivatives are included within the methods of the present invention. Analogs or derivatives are functional variants of an exendin or to
30 GLP-1 having similar amino acid sequence and retaining, to some extent, the increase in urine flow, increase in sodium excretion and/or decrease in potassium-excretion, activities

of the related exendin or GLP-1 or agonists thereto. By a functional variant is meant the derivative has an activity that can be substituted for one or more activities of a particular exendin or GLP-1 or an agonist thereto. Preferred functional variants retain all of the activities of a particular exendin or GLP-1 or an agonist thereto, however, the functional
5 variant may have an activity that, when measured quantitatively, is stronger or weaker, as measured in functional assays, for example, such as those disclosed herein. Preferred functional variants have activities that are within about 1% to about 10,000% of the activity of the related exendin, GLP-1, or agonist thereto, more preferably between about 10% to about 1000%, and more preferably within about 50% to about 500%. Derivatives
10 have at least about 50% sequence similarity, preferably about 70%, more preferably about 90%, and even more preferably about 95% sequence similarity to the related exendin or GLP-1, or agonist thereto. "Sequence similarity" refers to "homology" observed between amino acid sequences in two different polypeptides, irrespective of polypeptide origin.

The ability of the derivative to retain some activity can be measured using
15 techniques described herein. Derivatives include modification occurring during or after translation, for example, by phosphorylation, glycosylation, crosslinking, acylation, proteolytic cleavage, linkage to an antibody molecule, membrane molecule or other ligand (see Ferguson *et al.*, Annu. Rev. Biochem. 57:285-320, 1988).

Derivatives can be produced using standard chemical techniques and recombinant
20 nucleic acid molecule techniques. Modifications to a specific polypeptide may be deliberate, as through site-directed mutagenesis and amino acid substitution during solid-phase synthesis, or may be accidental such as through mutations in hosts which produce the polypeptide. Polypeptides including derivatives can be obtained using standard techniques such as those described in Sambrook, et al., *Molecular Cloning*, Cold Spring
25 Harbor Laboratory Press (1989).

The compounds referenced above form salts with various inorganic and organic acids and bases. Such salts include salts prepared with organic and inorganic acids, for example, HCl, HBr, H₂SO₄, H₃PO₄, trifluoroacetic acid, acetic acid, formic acid, methanesulfonic acid, toluenesulfonic acid, maleic acid, fumaric acid and camphorsulfonic
30 acid. Salts prepared with bases include ammonium salts, alkali metal salts, e.g. sodium and potassium salts, and alkali earth salts, e.g. calcium and magnesium salts. The salts

may be formed by conventional means, as by reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion
5 on a suitable ion exchange resin.

The claimed compositions can also be formulated as pharmaceutically acceptable salts (*e.g.*, acid addition salts) and/or complexes thereof. Pharmaceutically acceptable salts are non-toxic salts at the concentration at which they are administered. The preparation of such salts can facilitate the pharmacological use by altering the physical-chemical
10 characteristics of the composition without preventing the composition from exerting its physiological effect. Examples of useful alterations in physical properties include lowering the melting point to facilitate transmucosal administration and increasing the solubility to facilitate the administration of higher concentrations of the drug.

Pharmaceutically acceptable salts include acid addition salts such as those
15 containing sulfate, hydrochloride, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, cyclohexylsulfamate and quinate. Pharmaceutically acceptable salts can be obtained from acids such as hydrochloric acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic
20 acid, benzenesulfonic acid, *p*-toluenesulfonic acid, cyclohexylsulfamic acid, and quinic acid. Such salts may be prepared by, for example, reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion
25 on a suitable ion exchange resin.

The compounds described above are useful in view of their pharmacological properties. In particular, the compounds of the invention possess activity as agents to increase urine flow, increase sodium excretion and decrease potassium excretion, and to alleviate conditions or diseases associated with hypertoxic volemia.

30 Compositions useful in the invention may conveniently be provided in the form of formulations suitable for parenteral (including intravenous, intramuscular and

subcutaneous) or nasal or oral administration. In some cases, it will be convenient to provide an exendin or exendin agonist and another food-intake-reducing, plasma glucose-lowering or plasma lipid-lowering agent, such as amylin, an amylin agonist, a CCK, or a leptin, in a single composition or solution for administration together. In other cases, it may be more advantageous to administer the additional agent separately from said exendin or exendin agonist. A suitable administration format may best be determined by a medical practitioner for each patient individually. Suitable pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g., Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A.

10 "Parenteral Formulations of Proteins and Peptides: Stability and Stabilizers," Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2S (1988).

Compounds useful in the invention can be provided as parenteral compositions for injection or infusion. They can, for example, be suspended in an inert oil, suitably a vegetable oil such as sesame, peanut, olive oil, or other acceptable carrier. Preferably,

15 they are suspended in an aqueous carrier, for example, in an isotonic buffer solution at a pH of about 3.0 to 8.0, preferably at a pH of about 3.5 to 5.0. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH buffering agents. Useful buffers include

20 for example, sodium acetate/acetic acid buffers. A form of repository or "depot" slow release preparation may be used so that therapeutically effective amounts of the preparation are delivered into the bloodstream over many hours or days following transdermal injection or delivery.

The desired isotonicity may be accomplished using sodium chloride or other

25 pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol, polyols (such as mannitol and sorbitol), or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

Carriers or excipients can also be used to facilitate administration of the compound. Examples of carriers and excipients include calcium carbonate, calcium phosphate,

30 various sugars such as lactose, glucose, or sucrose, or types of starch, cellulose derivatives, gelatin, vegetable oils, polyethylene glycols and physiologically compatible solvents.

If desired, solutions of the above compositions may be thickened with a thickening agent such as methyl cellulose. They may be prepared in emulsified form, either water in oil or oil in water. Any of a wide variety of pharmaceutically acceptable emulsifying agents may be employed including, for example, acacia powder, a non-ionic surfactant (such as a Tween), or an ionic surfactant (such as alkali polyether alcohol sulfates or sulfonates, e.g., a Triton).

Compositions useful in the invention are prepared by mixing the ingredients following generally accepted procedures. For example, the selected components may be simply mixed in a blender or other standard device to produce a concentrated mixture which may then be adjusted to the final concentration and viscosity by the addition of water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity.

For use by the physician, the compositions will be provided in dosage unit form containing an amount of an exendin or exendin agonist, for example, exendin-3, and/or exendin-4. Therapeutically effective amounts of an exendin or exendin agonist for use in increasing urine flow are those that increase urine flow at a desired rate and level. As will be recognized by those in the field, an effective amount of therapeutic agent will vary with many factors including the age and weight of the patient, the patient's physical condition and other factors.

The effective dose of the compounds will typically be in the range of 1-30 μ g to about 10-20 mg, preferably about 30 μ g to 10 mg and more preferably about 300 μ g to 5 mg, most preferably 30 μ g to about 1 mg. The exact dose to be administered is determined by the attending clinician and is dependent, for example, upon where the particular compound lies within the above quoted range. Administration should begin whenever a diuretic effect is desired, for example, at the first sign of symptoms or shortly after diagnosis of renal failure, congestive heart failure, nephrotic syndrome, pulmonary edema, cirrhosis, hypertension, eclampsia, or pre-eclampsia. Administration may be by injection, preferably subcutaneous or intramuscular. Orally active compounds may be taken orally, however dosages should be increased 5-10 fold.

The optimal formulation and mode of administration of compounds of the present application to a patient depend on factors known in the art such as the particular disease or

disorder, the desired effect, and the type of patient. While the compounds will typically be used to treat human subjects they may also be used to treat similar or identical diseases in other vertebrates such as other primates, farm animals such as swine, cattle and poultry, and sports animals and pets such as horses, dogs and cats.

5 To assist in understanding the present invention, the following Examples are included. The experiments relating to this invention should not, of course, be construed as specifically limiting the invention and such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the invention as described herein and hereinafter claimed.

10

EXAMPLES

EXAMPLE 1: Diuretic Effects of GLP-1 or Exendin Administration

15 Materials: GLP-1 and exendin-4 were purchased from Bachem, Inc., Torrance, CA or synthesized at Amylin Pharmaceuticals, Inc., as described herein. Blood pressure transducers/transmitters were obtained from Data Sciences, Inc.

In Vivo Studies in Anesthetized rats: Male, Harlan Sprague Dawley rats were housed at 23±1°C in a 12:12 hour light:dark cycle (experiments being performed during
20 the light cycle) and were fed and watered *ad libitum* (Diet LM-485, Teklad, Madison, WI). Animals weighing 325-375 gms were fasted for ~20 hours prior to experimentation.

Surgical preparation: The preparation used here was as described in Young et al., (Drug Dev Res. 37:231-248, 1996), but modified by the addition of unilateral ureteral cannulation. Anesthesia was induced with 5% halothane, maintained with 2% halothane
25 during surgery and with 0.7 to 1% thereafter. Tracheotomy and cannulation of a femoral artery, saphenous vein and a single ureter were performed. The arterial line, perfused with heparinized saline (2U/ml), was used for blood sampling and pressure measurement (Spectramed P23XL transducer, Model 13-4615-58 amplifier, Gould, Cleveland, OH).
30 The venous line was used for drug administration. Total saline infusion rate was kept at 4 mL/hr. Colonic temperature was measured and controlled using a thermistor probe/controller (Model 73A, YSI, Yellow Springs, OH) and a heated operating table.

Signals for mean arterial pressure were periodically sampled at 1 Hz with 12 bit precision (DataTranslation DT2801A) and recorded (Labtech Notebook).

Numerical Methods: Dose-response curves were fitted to 4-parameter logistic functions and EC_{50} s derived using Prism (v2.0, GraphPad Software, San Diego, CA).

5 Observations are expressed as the percent of baseline, defined as the mean of measurements made in the 30 min prior to starting peptide or vehicle infusion. Data are expressed as mean \pm SEM. n=5-6.

Measurements: Samples of arterial blood (160 μ l) were collected periodically and samples of urine were collected every 15 min. Plasma and urinary sodium and potassium
10 concentrations were measured by ion-selective electrodes using Ciba/Corning 614 Na/K analyzer (Ciba/Corning, Inc., Medfield, MA). Unilateral urine flow was measured by weighing the 15-minute output of the cannulated ureter. Total urine flow was estimated as twice this amount.

Treatments: To obtain dose-responses, peptides were dissolved in 0.15 M NaCl
15 and administered as 0.1 ml bolus. Administration of GLP-1 had a strong effect on increasing urine flow ($ED_{50} = 0.71 \mu\text{g} \pm 0.26 \log \text{ units}$). The maximum response as percent of predose urine flow was $1764 \pm 281\%$ at 15 minutes for the $16.5 \mu\text{g}$ dose (Figures 3A-B). Administration of GLP-1 also increased sodium excretion (Figures 4A-B). However, GLP-1 significantly decreased the excretion of potassium (Figures 5A-B) ($ED_{50} \geq 0.25 \mu\text{g}$
20 $\pm 34 \log \text{ units}$ with a maximal fall to $13.9 \pm 1.7\%$ of predose concentrations at a dose of $1.65 \mu\text{g}$). Administration of exendin-4 also increased urine flow (Figures 8A-B). The ED_{50} was $0.12 \mu\text{g} \pm 0.18 \log \text{ units}$ and the maximum response as percent of predose urine flow was $2160 \pm 470\%$ at 15 minutes for the $21 \mu\text{g}$ dose. Administration of exendin also increased sodium excretion (Figures 9A-B). However, the excretion of potassium was
25 decreased (Figures 10A-B). The ED_{50} was $0.07 \mu\text{g} \pm 0.26 \log \text{ units}$ with a maximal decrease to $9.6 \pm 1.4\%$ of predose concentrations at a dose of $21 \mu\text{g}$.

EXAMPLE 2: Measurement of Arterial Blood Pressure and dP/dt in Conscious Rats by Telemetry After Administration of GLP-1 or Exendin-4:

30

Insertion of Transducers: Male, Harlan Sprague Dawley rats were anesthetized

with halothane and the abdominal aorta exposed after laparotomy. According to procedures detailed in the "Pressure Telemetry" manual from Data Sciences Inc., pressure transducer/transmitters were secured in place on the abdominal wall with the catheter tip in the abdominal aorta ~2mm above bifurcation. Following closure, the animals then
5 recovered to allow at least 7 days of stable recordings. Baseline data were collected during the 7+ days after surgery.

Measurement of blood pressure and dP/dt: After obtaining a stable baseline, rats received an intraperitoneal (ip) injection of GLP-1, exendin or vehicle alone (NaCl). The transmitted signals were recorded via telemetry and stored on a personal computer. Rate
10 of pressure change, dP/dt, was calculated by software provided by Data Sciences.

GLP-1: Animals received a single intraperitoneal (ip) injection of saline or GLP-1 (100 µl), n=7-8. Figures 1A-B depict the increase in mean arterial pressure after GLP-1 administration. Figure 2 depicts the increase in cardiac contractility after GLP-1 administration.

15 Exendin-4: Exendin-4 or saline (250 µl) was given twice daily (bid) by ip injection for five days. n=8 for saline and 5-6 for the exendin groups. Figures 6A-B depict the increase in mean arterial pressure after exendin-4 administration. Figure 7 depicts the increase in cardiac contractility after exendin-4 administration.

20 EXAMPLE 3

Cardiovascular actions of Exendin-4 or GLP-1 measured using transonic flow probes in anesthetised rats:

25 Materials, animal husbandry and cannulation under anesthesia: Materials animal husbandry and cannulation under anesthesia were as described in Example 1. Male Sprague Dawley rats (350-450g), anesthetised with halothane, were cannulated via the saphenous vein (for peptide injection) and femoral artery (for arterial pressure measurement).

30 Surgery: A transit time flow probe (2mm, 2SB, Transonic Systems Inc., Ithaca New York) was placed around the abdominal aorta, distal to renal, mesenteric and iliac

artery branches.

Measurements: The flow probe was connected to a Transonic TS-206 dual channel flowmeter for measurement of abdominal aortic blood flow. Heart rate was recorded using standard ECG electrodes. Peptides or vehicle (saline) were injected intravenously in a total volume of 100 μ L over 1-2 minutes. Mean arterial pressure (MAP), heart rate (HR) and mean aortic blood flow, were recorded every second using Labtek Notebook data acquisition software over the experimental period. Aortic conductance (flow/MAP; mL/min/mmHg) and stroke volume (flow/HR; mL/min per beats/min = mL) were then derived.

Treatments: Exendin-4 was injected in doses of 0.021, 0.21, 2.1 and 21 μ g, and GLP-1 was injected in doses of 0.0165, 0.165, 1.65 and 16.5 μ g after a 20 minute control period.

Results:

GLP-1: GLP-1 at a dose of 16.5 μ g increased mean arterial pressure by 22mmHg within 5 minutes of administration. Aortic blood flow increased by 57% from 14 to 22 mL/min, heart rate by 17% from 360 to 420 beat/min, stroke volume by 38% from 37 to 51 μ L, and aortic conductance by 50% from 0.12 to 0.18 mL/min/mmHg within 2 minutes of GLP-1 administration. Effects lasted for about 10 min.

Exendin-4: A similar pattern of effects was observed with a 0.21 μ g dose of exendin-4 (~30mmHg increase in blood pressure; 60% increase in aortic blood flow; 40% increase in heart rate; 60% increase in stroke volume; 35% increase in aortic conductance), except that effects persisted for 30-60 min. These responses, where there are large changes in aortic blood flow and lesser changes in blood pressure, are consistent with GLP-1 and exendin-4 having inotropic (cardiac stimulatory) and vasodilator properties.

25

EXAMPLE 4Preparation of Peptide having SEQ. ID. NO. 5

5 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly-NH₂ [SEQ. ID. NO. 5]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g)
10 using Fmoc-protected amino acids (Applied Biosystems, Inc.). In general, single-coupling
cycles were used throughout the synthesis and Fast Moc (HBTU activation) chemistry was
employed. Deprotection (Fmoc group removal) of the growing peptide chain was achieved
using piperidine. Final deprotection of the completed peptide resin was achieved using a
mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2
15 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to
Cleavage Techniques, Applied Biosystems, Inc.) The peptide was precipitated in
ether/water (50 mL) and centrifuged. The precipitate was reconstituted in glacial acetic
acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was
about 75%.

20 Used in purification steps and analysis were Solvent A (0.1% TFA in water) and
Solvent B (0.1% TFA in ACN).

The solution containing peptide was applied to a preparative C-18 column and purified
(10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined
isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the
25 above-identified peptide. Analytical RP-HPLC (gradient 30% to 50% Solvent B in Solvent
A over 30 minutes) of the lyophilized peptide gave product peptide having an observed
retention time of 18.9 minutes. Electrospray Mass Spectrometry (M): calculated 3408.0;
found 3408.9.

30 EXAMPLE 5

Preparation of Peptide having SEQ. ID. NO. 6

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ. ID. NO. 6]

5

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g)
using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin,
deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A
10 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30%
to 40% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 17.9 minutes. Electrospray Mass Spectrometry
(M): calculated 3294.7; found 3294.8.

15 EXAMPLE 6Preparation of Peptide having SEQ. ID. NO. 7

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
20 Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 7]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
25 resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
29% to 36% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 20.7 minutes. Electrospray Mass Spectrometry
(M): calculated 3237.6; found 3240.

30

EXAMPLE 7

Preparation of Peptide having SEQ. ID. NO. 8

His Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
5 Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 8]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g)
using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin,
10 deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36%
to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 15.2 minutes. Electrospray Mass Spectrometry
(M): calculated 3251.6; found 3251.5.

15

EXAMPLE 8

Preparation of Peptide having SEQ. ID. NO. 9

20 His Gly Glu Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val
Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 9]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g)
25 using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin,
deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36%
to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 13.1 minutes. Electrospray Mass Spectrometry
30 (M): calculated 3207.6; found 3208.3.

EXAMPLE 9Preparation of Peptide having SEQ. ID. NO. 10

5 His Gly Glu Gly Thr Ala Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 10]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g)
10 using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin,
deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 35%
to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 12.8 minutes. Electrospray Mass Spectrometry
15 (M): calculated 3161.5; found 3163.

EXAMPLE 10Preparation of Peptide having SEQ. ID. NO. 11

20

His Gly Glu Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 11]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
25 Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
30 peptide having an observed retention time of 15.2 minutes. Electrospray Mass Spectrometry
(M): calculated 3221.6; found 3222.7.

EXAMPLE 11Preparation of Peptide having SEQ. ID. NO. 12

5

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 12]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
10 Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
34% to 44% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
15 peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry
(M): calculated 3195.5; found 3199.4.

EXAMPLE 12

20

Preparation of Peptide having SEQ. ID. NO. 13

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 13]

25 The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 15.7 minutes. Electrospray Mass Spectrometry

(M): calculated 3221.6; found 3221.6.

EXAMPLE 13

5 Preparation of Peptide having SEQ. ID. NO. 14

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 14]

10 The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
15 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 18.1 minutes. Electrospray Mass Spectrometry
(M): calculated 3180.5; found 3180.9.

EXAMPLE 14

20

Preparation of Peptide having SEQ. ID. NO. 15

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 15]

25

 The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
30 A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product

peptide having an observed retention time of 17.0 minutes. Electrospray Mass Spectrometry (M): calculated 3180.6; found 3182.8.

EXAMPLE 15

5

Preparation of Peptide having SEQ. ID. NO. 16

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 16]

10

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
15 A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.9 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3195.9.

20 EXAMPLE 16

Preparation of Peptide having SEQ. ID. NO. 17

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Ala Glu Glu Ala Val Arg Leu Phe
25 Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 17]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
30 resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient

37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3179.0.

5 EXAMPLE 17

Preparation of Peptide having SEQ. ID. NO. 18

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Ala Glu Ala Val Arg Leu Phe
10 Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 18]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
15 resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.0.

20

EXAMPLE 18

Preparation of Peptide having SEQ. ID. NO. 19

25 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Ala Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 19]

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g)
30 using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A

(0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 13.7 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3179.0.

5

EXAMPLE 19

Preparation of Peptide having SEQ. ID. NO. 20

10 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Ala Arg Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 20]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
35% to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 14.0 minutes. Electrospray Mass Spectrometry
20 (M): calculated 3209.6; found 3212.8.

EXAMPLE 20

Preparation of Peptide having SEQ. ID. NO. 21

25

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Ala Leu
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 21]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
30 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the

resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3152.5; found 3153.5.

EXAMPLE 21

Preparation of Peptide having SEQ. ID. NO. 22

10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Ala
Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 22]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
35% to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product
peptide having an observed retention time of 12.1 minutes. Electrospray Mass Spectrometry
(M): calculated 3195.5; found 3197.7.

EXAMPLE 22

25

Preparation of Peptide having SEQ. ID. NO. 23

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Ala Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 23]

30

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55

mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 10.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.5.

EXAMPLE 23

10 Preparation of Peptide having SEQ. ID. NO. 24

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Ala Leu Lys Asn-NH₂ [SEQ. ID. NO. 24]

15 The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 20 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.5 minutes. Electrospray Mass Spectrometry (M): calculated 3161.5; found 3163.0.

EXAMPLE 24

25

Preparation of Peptide having SEQ. ID. NO. 25

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Ala Lys Asn-NH₂ [SEQ. ID. NO. 25]

30

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-

Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.5 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3199.

EXAMPLE 25

Preparation of Peptide having SEQ. ID. NO. 26

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Ala Asn-NH₂ [SEQ. ID. NO. 26]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.5 minutes. Electrospray Mass Spectrometry (M): calculated 3180.5; found 3183.7.

EXAMPLE 26

Preparation of Peptide having SEQ. ID. NO. 27

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Ala-NH₂ [SEQ. ID. NO. 27]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 34% to 44% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 22.8 minutes. Electrospray Mass Spectrometry (M): calculated 3194.6; found 3197.6.

10 EXAMPLE 27

Preparation of Peptide having SEQ. ID. NO. 28

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
15 Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID.
NO. 28]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
20 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass
25 Spectrometry (M): calculated 4099.6.

EXAMPLE 28

Preparation of Peptide having SEQ. ID. NO. 29

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
5 Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID.
NO. 29]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
10 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then
carried out to determine the retention time of the product peptide. Electrospray Mass
15 Spectrometry (M): calculated 4042.5.

EXAMPLE 29Preparation of Peptide having SEQ. ID. NO. 30

20

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO.
30]

25 The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g)
using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin,
deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1%
TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to
30 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out
to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 4002.4

EXAMPLE 30

5 Preparation of Peptide having SEQ. ID. NO. 31

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro- NH_2 [SEQ. ID. NO.
31]

10

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent
15 A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then
carried out to determine the retention time of the product peptide. Electrospray Mass
Spectrometry (M): calculated 3945.4.

20 EXAMPLE 31

Preparation of Peptide having SEQ. ID. NO. 32

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val
25 Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro- NH_2 [SEQ. ID.
NO. 32]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
30 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent

A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3905.3.

5

EXAMPLE 32

Preparation of Peptide having SEQ. ID. NO. 33

10 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro-NH₂ [SEQ. ID. NO. 33]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass
20 Spectrometry (M): calculated 3848.2.

EXAMPLE 33

Preparation of Peptide having SEQ. ID. NO. 34

25

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 34]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
30 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the

resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3808.2.

EXAMPLE 34

Preparation of Peptide having SEQ. ID. NO. 35

10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 35]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then
carried out to determine the retention time of the product peptide. Electrospray Mass
Spectrometry (M): calculated 3751.1.

EXAMPLE 35

25

Preparation of Peptide having SEQ. ID. NO. 36

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly-NH₂ [SEQ. ID. NO. 36]

30

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55

mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3737.1.

EXAMPLE 36

Preparation of Peptide having SEQ. ID. NO. 37

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser Gly-NH₂ [SEQ. ID. NO. 37]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1.

EXAMPLE 37

25

Preparation of Peptide having SEQ. ID. NO. 38

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser-NH₂ [SEQ. ID. NO. 38]

30

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-

Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
5 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1

EXAMPLE 38

10

Preparation of Peptide having SEQ. ID. NO. 39

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser-NH₂ [SEQ. ID. NO. 39]

15

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent
20 A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3623.0.

25 EXAMPLE 39

Preparation of Peptide having SEQ. ID. NO. 40

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
30 Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser-NH₂ [SEQ. ID. NO. 40]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3593.0

10 EXAMPLE 40

Preparation of Peptide having SEQ. ID. NO. 41

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
15 Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser-NH₂ [SEQ. ID. NO. 41]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3535.9

25

EXAMPLE 41

Preparation of Peptide having SEQ. ID. NO. 42

30 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro-NH₂ [SEQ. ID. NO. 42]

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3505.9.

10

EXAMPLE 42Preparation of Peptide having SEQ. ID. NO. 43

15 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro-NH₂ [SEQ. ID. NO. 43]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3448.8.

25

EXAMPLE 43Preparation of Peptide having SEQ. ID. NO. 44

30

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu

Phe Ile Glu Phe Leu Lys Asn Gly Gly-NH₂ [SEQ. ID. NO. 44]

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3351.7.

EXAMPLE 44

. Preparation of Peptide having SEQ. ID. NO. 45

15

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly-NH₂ [SEQ. ID. NO. 45]

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3351.8.

EXAMPLE 45

30

Preparation of Peptide having SEQ. ID. NO. 46

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly-NH₂ [SEQ. ID. NO. 46]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
5 Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then
10 carried out to determine the retention time of the product peptide. Electrospray Mass
Spectrometry (M): calculated 3294.7.

EXAMPLE 46

15 Preparation of Peptide having SEQ. ID. NO. 47

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly tPro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID.
NO. 47]

20 The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Double couplings are required
25 at residues 37,36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B
(0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A
over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time
of the product peptide. Electrospray Mass Spectrometry (M): calculated 4197.1.

30 EXAMPLE 47

Preparation of Peptide having SEQ. ID. NO. 48

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID.
5 NO. 48]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
10 resin, deprotected and purified in a similar way to Example 4. Double couplings are required
at residues 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent
B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A
over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time
of the product peptide. Electrospray Mass Spectrometry (M): calculated 4179.1.

15

EXAMPLE 48Preparation of Peptide having SEQ. ID. NO. 49

20 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly NMeala Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID.
NO. 49]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
25 Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Double couplings are required
at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B
(0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A
30 over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time
of the product peptide. Electrospray Mass Spectrometry (M): calculated 3948.3.

EXAMPLE 49Preparation of Peptide having SEQ. ID. NO. 50

5

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly NMeala Ser Ser Gly Ala NMeala Nmeala-*NH*₂
[SEQ. ID. NO. 50]

10 The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Double couplings are required
at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B
15 (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A
over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time
of the product peptide. Electrospray Mass Spectrometry (M): calculated 3840.1.

EXAMPLE 50

20

Preparation of Peptide having SEQ. ID. NO. 51

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly hPro Ser Ser Gly Ala hPro hPro-*NH*₂ [SEQ. ID.
25 NO. 51]

30 The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Double couplings are required
at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B

(0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4050.1.

5 EXAMPLE 51

Preparation of Peptide having SEQ. ID. NO. 52

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
10 Phe Ile Glu Trp Leu Lys Asn Gly Gly hPro Ser Ser Gly Ala hPro-NH₂ [SEQ. ID. NO. 52]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
15 resin, deprotected and purified in a similar way to Example 4. A double coupling is required at residue 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3937.1

20

EXAMPLE 52

Preparation of Peptide having SEQ. ID. NO. 53

25 Arg Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 53]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
30 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent

A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3827.2.

5

EXAMPLE 53

Preparation of Peptide having SEQ. ID. NO. 54

10 His Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly Gly-NH₂ [SEQ. ID. NO. 54]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass
20 Spectrometry (M): calculated 3394.8.

EXAMPLE 54

Preparation of Peptide having SEQ. ID. NO. 55

25

His Gly Glu Gly Thr Naphthylala Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val
Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 55]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
30 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the

resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3289.5.

EXAMPLE 55

Preparation of Peptide having SEQ. ID. NO. 56

10

His Gly Glu Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ. ID. NO. 56]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the
resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then
carried out to determine the retention time of the product peptide. Electrospray Mass
Spectrometry (M): calculated 3280.7.

EXAMPLE 56

25

Preparation of Peptide having SEQ. ID. NO. 57

His Gly Glu Gly Thr Phe Ser Thr Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ. ID. NO. 57]

30

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55

mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then
5 carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3294.7.

EXAMPLE 57

10 Preparation of Peptide having SEQ. ID. NO. 58

His Gly Glu Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Met Ala Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ. ID. NO. 58]

15 The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
20 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3250.7.

EXAMPLE 58

25

Preparation of Peptide having SEQ. ID. NO. 59

His Gly Glu Gly Thr Phe Thr Ser Asp pentylgly Ser Lys Gln Leu Glu Glu Glu Ala Val
Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 59]

30

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-

Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3253.5.

EXAMPLE 59

Preparation of Peptide having SEQ. ID. NO. 60

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 60]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3289.5.

EXAMPLE 60

Preparation of Peptide having SEQ. ID. NO. 61

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe tButylgly Glu Trp Leu Lys Asn-NH₂ [SEQ. ID. NO. 61]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3183.4.

10 EXAMPLE 61

Preparation of Peptide having SEQ. ID. NO. 62

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
15 Phe Ile Asp Phe Leu Lys Asn-NH₂ [SEQ. ID. NO. 62]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3237.6.

25

EXAMPLE 62

Preparation of Peptide having SEQ. ID. NO. 63

30 His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser-NH₂ [SEQ. ID. NO. 63]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3637.9.

10

EXAMPLE 63Preparation of Peptide having SEQ. ID. NO. 64

15 His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys Asn Gly-NH₂ [SEQ. ID. NO. 64]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3309.7.

25

EXAMPLE 64Preparation of Peptide having SEQ. ID. NO. 65

30

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu

Phe Ile Glu Trp Leu Lys Asn Gly Gly hPro Ser Ser Gly Ala hPro hPro-NH₂ [SEQ. ID. NO. 65]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-
5 Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 4. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A
10 over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3711.1.

EXAMPLE 65

15 Preparation of C-terminal carboxylic acid peptides corresponding to the above C-terminal amide sequences for SEQ. ID. NOS. 5-27, 34-41, 44-46 and 53-64

Peptides having the sequences of SEQ. ID. NOS. 5-27, 34-41, 44-46 and 53-64 are assembled on the so called Wang resin (p-alkoxybenzylalcohol resin (Bachem, 0.54
20 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass
25 Spectrometry provides an experimentally determined (M).

EXAMPLE 66

Preparation of C-terminal carboxylic acid peptides corresponding to the above C-terminal
30 amide sequences for SEQ. ID. NOS. 28-33, 42, 43, 47-52 and 65

Peptides having the sequence of SEQ. ID. NOS. 28-33, 42, 43, 47-52 and 65 are assembled on the 2-chlorotritylchloride resin (200-400 mesh), 2% DVB (Novabiochem, 0.4-1.0 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are

5 Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

WE CLAIM:

1. A method for increasing urine flow in an individual comprising administering a therapeutically effective amount of an exendin or an exendin agonist.

5

2. The method of claim 1, wherein said increase in urine flow is accompanied by an increase in sodium excretion in said individual.

3. The method of claim 1, wherein said increase in urine flow does not increase urinary potassium concentration in said individual.

10

4. A method of decreasing the concentration of potassium in the urine of an individual comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

15

5. A method of preventing or alleviating a condition or disorder associated with toxic hypervolemia in an individual, comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

6. The method of claim 5, wherein condition or disorder is selected from the group consisting of renal failure, congestive heart failure, nephrotic syndrome, pulmonary edema and cirrhosis.

20

7. The method of claim 5, wherein said condition or disorder is hypertension.

25

8. A method of inducing rapid diuresis in an individual comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

9. A method of preparing an individual for a surgical procedure comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

30

10. The method of claim 9, wherein said surgical procedure is selected from the group consisting of ocular surgical procedures and neurosurgical procedures.

5 11. The method of claim 9, wherein said exendin or exendin agonist is administered to said individual before said surgical procedure.

12. A method of increasing renal plasma flow and glomerular filtration rate in an individual comprising administering to said individual a therapeutically effective amount of
10 an exendin or an exendin agonist.

13. A method of treating pre-eclampsia or eclampsia of pregnancy in an individual comprising administering to said individual a therapeutically effective amount of an exendin or an exendin agonist.

15

14. The method according to any of claims 1, 4, 5, 8, 9, 12, or 13, wherein said exendin is exendin-3 [SEQ ID NO. 1].

15. The method according to any of claims 1, 4, 5, 8, 9, 12, or 13, wherein said
20 exendin is exendin-4 [SEQ ID NO. 2].

16. The method according to any of claims 1, 4, 5, 8, 9, 12 or 13, wherein said exendin agonist is a peptide compound of the formula [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
25 Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

Xaa₁ is His, Arg or Tyr;

Xaa₂ is Ser, Gly, Ala or Thr;

30 Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

- Xaa₆ is Ala, Phe, Tyr or naphthylalanine;
Xaa₇ is Thr or Ser;
Xaa₈ is Ala, Ser or Thr;
Xaa₉ is Asp or Glu;
5 Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
Xaa₁₁ is Ala or Ser;
Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
10 Xaa₁₅ is Ala or Glu;
Xaa₁₆ is Ala or Glu;
Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
15 Xaa₂₁ is Ala or Leu;
Xaa₂₂ is Phe, Tyr or naphthylalanine;
Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
or Met;
Xaa₂₄ is Ala, Glu or Asp;
20 Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
Xaa₂₆ is Ala or Leu;
Xaa₂₇ is Ala or Lys;
Xaa₂₈ is Ala or Asn;
Z₁ is -OH,
25 -NH₂,
Gly-Z₂,
Gly Gly-Z₂,
Gly Gly Xaa₃₁-Z₂,
Gly Gly Xaa₃₁ Ser-Z₂,
30 Gly Gly Xaa₃₁ Ser Ser-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂, or
 5 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈ Xaa₃₉-Z₂; wherein Xaa₃₁,
 Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently
 selected from the group consisting of Pro,
 homoproline, 3Hyp, 4Hyp, thioproline,
 N-alkylglycine, N-alkylpentylglycine and
 10 N-alkylalanine; Xaa₃₉ is Ser, Thr or Tyr; and
 Z₂ is -OH or -NH₂; and pharmaceutically acceptable salts thereof;
 provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃,
 Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇, and Xaa₂₈ are
 Ala; and provided also that the compound is not exendin-3 [SEQ ID NO. 1] or exendin-4
 15 [SEQ ID NO. 2].

17. The method according to any of claims 1, 4, 5, 8, 9, 12, or 13, wherein said exendin or exendin agonist is administered peripherally.
- 20 18. The method according to claim 17, wherein said exendin or exendin agonist is administered subcutaneously.
19. The method of claim 17, wherein said peripheral administration is selected from the group consisting of buccal, nasal, pulmonary, oral, intravenous, intraocular,
 25 rectal, and transdermal administration.
20. A pharmaceutical composition for use in the treatment of conditions or disorders associated with hypervolemia comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.
- 30 21. A pharmaceutical composition for use in increasing urine flow in an

individual comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

22. A pharmaceutical composition for use in treating pre-eclampsia or
5 eclampsia of pregnancy in an individual comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

23. A pharmaceutical composition according to any of claims 20, 21 or 22,
10 wherein said exendin is exendin-3 [SEQ ID NO. 1].

24. A pharmaceutical composition according to any of claims 20, 21 or 22,
wherein said exendin is exendin-4 [SEQ ID NO. 2].

25. A pharmaceutical composition according to any of claims 22, 23 or 24,
15 wherein said exendin agonist is a peptide compound of the formula [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

20 Xaa₁ is His, Arg or Tyr;
Xaa₂ is Ser, Gly, Ala or Thr;
Xaa₃ is Asp or Glu;
Xaa₅ is Ala or Thr;
Xaa₆ is Ala, Phe, Tyr or naphthylalanine;
25 Xaa₇ is Thr or Ser;
Xaa₈ is Ala, Ser or Thr;
Xaa₉ is Asp or Glu;
Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
Xaa₁₁ is Ala or Ser;
30 Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;

- Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
 Xaa₁₅ is Ala or Glu;
 Xaa₁₆ is Ala or Glu;
 Xaa₁₇ is Ala or Glu;
 5 Xaa₁₉ is Ala or Val;
 Xaa₂₀ is Ala or Arg;
 Xaa₂₁ is Ala or Leu;
 Xaa₂₂ is Phe, Tyr or naphthylalanine;
 Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
 10 or Met;
 Xaa₂₄ is Ala, Glu or Asp;
 Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
 Xaa₂₆ is Ala or Leu;
 Xaa₂₇ is Ala or Lys;
 15 Xaa₂₈ is Ala or Asn;
 Z₁ is -OH,
 -NH₂,
 Gly-Z₂,
 Gly Gly-Z₂,
 20 Gly Gly Xaa₃₁-Z₂,
 Gly Gly Xaa₃₁ Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
 25 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂, or
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈ Xaa₃₉-Z₂;
 wherein Xaa₃₁, Xaa₃₆, Xaa₃₇, and Xaa₃₈ are independently
 30 selected from the group consisting of Pro,
 homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine and

N-alkylalanine; Xaa₃₉ is Ser, Thr or Tyr; and

Z₂ is -OH or -NH₂; and pharmaceutically acceptable salts thereof;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃,

- 5 Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇, and Xaa₂₈ are Ala; and provided also that the compound is not exendin-3 [SEQ ID NO. 1] or exendin-4 [SEQ ID NO. 2].

26. A method of increasing urine flow in an individual comprising
10 administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist.

27. The method of claim 26, wherein said increase in urine flow is
accompanied by an increase in sodium excretion in said individual.

15

28. The method of claim 26, wherein said increase in urine flow does not
increase urinary potassium concentration in said individual.

29. A method of decreasing the concentration of potassium in the urine of an
20 individual comprising administering to said individual a therapeutically effective amount
of GLP-1 or a GLP-1 agonist.

30. A method of preventing or alleviating a condition or disorder associated
with toxic hypervolemia in an individual, comprising administering to said individual a
25 therapeutically effective amount of GLP-1 or a GLP-1 agonist.

31. The method of claim 30, wherein said condition or disorder is selected from
the group consisting of renal failure, congestive heart failure, nephrotic syndrome,
pulmonary edema and cirrhosis.

30

32. The method of claim 30, wherein said condition or disorder is hypertension.

33. A method of inducing rapid diuresis in an individual comprising administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist.

5

34. A method of preparing an individual for a surgical procedure comprising administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist.

10 35. The method of claim 34, wherein said surgical procedure is selected from the group consisting of ocular surgical procedures and neurosurgical procedures.

36. The method of claim 34, wherein said GLP-1 or a GLP-1 agonist is administered to said individual before said surgical procedure.

15

37. A method of increasing renal plasma flow and glomerular filtration rate in an individual comprising administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist.

20 38. A method of treating pre-eclampsia or eclampsia of pregnancy in an individual comprising administering to said individual a therapeutically effective amount of GLP-1 or a GLP-1 agonist.

25 39. The method according to any of claims 26, 29, 30, 33, 34, 37 or 38, wherein said GLP-1 or GLP-1 agonist is administered peripherally.

40. The method of claim 39, wherein said GLP-1 or a GLP-1 agonist is administered subcutaneously.

30 41. The method of claim 39, wherein said peripheral administration is selected from the group consisting of buccal, nasal, pulmonary, oral, intravenous, intraocular,

rectal, and transdermal administration.

42. A pharmaceutical composition for use in the treatment of conditions or disorders associated with hypervolemia comprising a therapeutically effective amount of GLP-1 or a GLP-1 agonist in association with a pharmaceutically acceptable carrier.

43. A pharmaceutical composition for use in increasing urine flow in an individual comprising a therapeutically effective amount of GLP-1 or a GLP-1 agonist in association with a pharmaceutically acceptable carrier.

44. A pharmaceutical composition for use in treating pre-eclampsia or eclampsia of pregnancy in an individual comprising a therapeutically effective amount of GLP-1 or a GLP-1 agonist in association with a pharmaceutically acceptable carrier.

45. A method for increasing cardiac contractility in an individual comprising administering a therapeutically effective amount of an exendin or an exendin agonist.


46. A method for treating a condition or disorder that can be alleviated by increasing cardiac contractility in an individual comprising administering a therapeutically effective amount of an exendin or an exendin agonist.

47. The method according to claim 46 wherein said condition or disorder is selected from the group consisting of congestive heart failure, pulmonary edema, systemic edema, and renal failure.

48. The method according to claim 45 or claim 46 wherein said exendin is exendin-3 [SEQ ID NO. 1].

49. The method according to claim 45 or claim 46 wherein said exendin is exendin-4 [SEQ ID NO. 2].

50. The method according to claim 45 or 46 wherein said exendin agonist is a peptide compound of the formula [SEQ. ID. NO. 4]:



- Xaa₁ is His, Arg or Tyr;
Xaa₂ is Ser, Gly, Ala or Thr;
Xaa₃ is Asp or Glu;
10 Xaa₅ is Ala or Thr;
Xaa₆ is Ala, Phe, Tyr or naphthylalanine;
Xaa₇ is Thr or Ser;
Xaa₈ is Ala, Ser or Thr;
Xaa₉ is Asp or Glu;
15 Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
Xaa₁₁ is Ala or Ser;
Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
20 Xaa₁₅ is Ala or Glu;
Xaa₁₆ is Ala or Glu;
Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
25 Xaa₂₁ is Ala or Leu;
Xaa₂₂ is Phe, Tyr or naphthylalanine;
Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
or Met;
Xaa₂₄ is Ala, Glu or Asp;
30 Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
Xaa₂₆ is Ala or Leu;

Xaa₂₇ is Ala or Lys;

Xaa₂₈ is Ala or Asn;

Z₁ is -OH,

-NH₂,

5

Gly-Z₂,

Gly Gly-Z₂,

Gly Gly Xaa₃₁-Z₂,

Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,

10

Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂, or

15

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈ Xaa₃₉-Z₂;

wherein Xaa₃₁, Xaa₃₆, Xaa₃₇, and Xaa₃₈ are independently

selected from the group consisting of Pro,

homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine and

20

N-alkylalanine; Xaa₃₉ is Ser, Thr or Tyr; and

Z₂ is -OH or -NH₂; and pharmaceutically acceptable salts thereof;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃,

Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇, and Xaa₂₈ are

Ala; and provided also that the compound is not exendin-3 [SEQ ID NO. 1] or exendin-4

25

[SEQ ID NO. 2].

51. The method according to claim 45 or claim 46 wherein said exendin or exendin agonist is administered peripherally.

30

52. The method according to claim 51 wherein said peripheral administration is selected from the group consisting of buccal, nasal, pulmonary, oral, intraocular, rectal,

and transdermal administration.

53. The method according to claim 45 or claim 46 wherein said exendin or exendin agonist is administered intravenously.

5

54. A pharmaceutical composition for use in the treatment of a condition or disorder that can be alleviated by increasing cardiac contractility comprising a therapeutically effective amount of an exendin or an exendin agonist in association with a pharmaceutically acceptable carrier.

10

55. The pharmaceutical composition of claim 54 wherein said exendin is exendin-3 [SEQ ID NO. 1].

56. The pharmaceutical composition of claim 54 wherein said exendin is exendin-4 [SEQ. ID NO. 2].

15

57. The pharmaceutical composition of claim 54 wherein said exendin agonist is a peptide compound of the formula [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

20

Xaa₁ is His, Arg or Tyr;

Xaa₂ is Ser, Gly, Ala or Thr;

25 Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

Xaa₆ is Ala, Phe, Tyr or naphthylalanine;

Xaa₇ is Thr or Ser;

Xaa₈ is Ala, Ser or Thr;

30 Xaa₉ is Asp or Glu;

Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;

- Xaa₁₁ is Ala or Ser;
 Xaa₁₂ is Ala or Lys;
 Xaa₁₃ is Ala or Gln;
 Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
 5 Xaa₁₅ is Ala or Glu;
 Xaa₁₆ is Ala or Glu;
 Xaa₁₇ is Ala or Glu;
 Xaa₁₉ is Ala or Val;
 Xaa₂₀ is Ala or Arg;
 10 Xaa₂₁ is Ala or Leu;
 Xaa₂₂ is Phe, Tyr or naphthylalanine;
 Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
 or Met;
 Xaa₂₄ is Ala, Glu or Asp;
 15 Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
 Xaa₂₆ is Ala or Leu;
 Xaa₂₇ is Ala or Lys;
 Xaa₂₈ is Ala or Asn;
 Z₁ is -OH,
 20 -NH₂,
 Gly-Z₂,
 Gly Gly-Z₂,
 Gly Gly Xaa₃₁-Z₂,
 Gly Gly Xaa₃₁ Ser-Z₂,
 25 Gly Gly Xaa₃₁ Ser Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂,
 30 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂, or
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈ Xaa₃₉-Z₂;

wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from the group consisting of Pro, homoproline, 3Hyp, 4Hyp, thioproline, N-alkylglycine, N-alkylpentylglycine and N-alkylalanine; Xaa₃₉ is Ser, Thr or Tyr; and Z₂ is -OH or -NH₂; and pharmaceutically acceptable salts thereof; provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇, and Xaa₂₈ are Ala; and provided also that the compound is not exendin-3 [SEQ ID NO. 1] or exendin-4 [SEQ ID NO. 2].

58. A method for increasing cardiac contractility in an individual comprising administering a therapeutically effective amount of GLP-1 or a GLP-1 agonist.

59. A method for treating a condition or disorder that can be alleviated by increasing cardiac contractility in an individual comprising administering a therapeutically effective amount of a GLP-1 or a GLP-1 agonist.

60. The method of claim 59 wherein said condition or disorder is selected from the group consisting of congestive heart failure, pulmonary edema, systemic edema, and renal failure.

61. The method according to claim 58 or claim 59 wherein said GLP-1 or GLP-1 agonist is administered peripherally.

62. The method according to claim 61 wherein said peripheral administration is selected from the group consisting of buccal, nasal, pulmonary, oral, intraocular, rectal, and transdermal administration.

63. The method according to claim 58 or claim 59 wherein said GLP-1 or GLP-1 agonist is administered intravenously.

64. A pharmaceutical composition for use in the treatment of a condition or disorder that can be alleviated by increasing cardiac contractility comprising a therapeutically effective amount of a GLP-1 or a GLP-1 agonist in association with a
- 5 pharmaceutically acceptable carrier.

1/10

RESPONSE OF MAP TO GLP-1




FIG. 1A





DOSE-RESPONSE CURVE:
MAP TO GLP-1

FIG. 1B



2/10

*FIG. 2*


3/10

*FIG. 3A**FIG. 3B*


4/10

*FIG. 4A**FIG. 4B*

5/10

*FIG. 5A*

DOSE-RESPONSE CURVE:
URINE POTASSIUM CONCENTRATION TO GLP-1

*FIG. 5B*

6/10

RESPONSE OF MAP TO EXENDIN-4




FIG. 6A



DOSE-RESPONSE CURVE:
MAP TO EXENDIN-4

FIG. 6B

7/10

*FIG. 7*

8/10

RESPONSE OF URINE FLOW TO EXENDIN-4




FIG. 8A


DOSE-RESPONSE CURVE:
URINE FLOW TO EXENDIN-4

FIG. 8B

9/10

RESPONSE OF SODIUM EXCRETION TO EXENDIN-4




FIG. 9A




DOSE-RESPONSE CURVE:
SODIUM EXCRETION TO EXENDIN-4

FIG. 9B

10/10

RESPONSE OF URINE POTASSIUM CONCENTRATION TO EXENDIN-4

*FIG. 10A**FIG. 10B*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/02554

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A01N 37/18
US CL :514/2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,512,549 A (CHEN et al) 30 April 1996, see entire document.	1-23, 25-64
Y	US 5,424,286 A (ENG) 13 June 1995, see entire document.	1-23, 25-64

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 MARCH 1999

Date of mailing of the international search report

13 APR 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized Officer

L. BLAINE LANKFORD

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/02554

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 24
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.